





TREATMENT CONSIDERATIONS FORM

INSTRUCTIONS: Please read and initial after each paragraph to acknowledge that you understand the terms of this treatment, unless told otherwise.

The CoolSculpting® procedure is a non-invasive procedure that is intended to change the appearance of the treatment area

by delivering controlled cooling at the surface of the skin to break down fat cells that are just beneath the skin. This procedure is not a treatment for obesity or a weight-loss solution. The CoolSculpting procedure does not replace traditional methods such as diet, exercise or liposuction. Initial:
Clinical studies of a treatment site have shown that the CoolSculpting procedure can break down fat cells to change the appearance of visibly localized bulges of fat that is just beneath the skin on the abdomen, thighs, flanks and submental area. The submental area is the area under the chin. Following the procedure, the treated fat cells are naturally processed by the body. Visible results can vary from person to person. Initial:
WHAT YOU CAN EXPECT:
Temporary Sensations/Symptoms:
» The suction pressure of a vacuum applicator may cause sensations of deep pulling, tugging and pinching. A surface applicator may cause sensations of pressure. You may experience intense cold, stinging, tingling, aching or cramping as the treatment begins. These sensations generally subside during treatment as the area becomes numb. Initial:
» You may have dizziness, light-headedness, nausea, flushing, sweating, or fainting during or immediately after the treatment Initial :
»The treated area may look or feel stiff after the procedure and transient blanching (temporary whitening of the skin) may occur. These are all normal reactions that typically resolve within a few minutes. Initial:
» Bruising, swelling, redness, cramping and pain can occur in the treated area and the treated area may appear red for one to two weeks after treatment. Initial:
» After submental area treatment, a feeling of fullness in the back of the throat may occur. You should initial here only if the

Page 1 of 3

submental area is to be treated. If the area under the chin is not being treated, please write N/A. Initial:

» You may feel a dulling of sensation in the treated area that can last for several weeks after the procedure. Prolonged swelling, itching, tingling, numbness, tenderness to the touch, pain in the treated area, cramping, aching, bruising and/or ski sensitivity also have been reported. Initial:
Potential Side Effects / Risks
»Paradoxical Hyperplasia A small number of patients have experienced gradual development of a firmer enlargement of varying size and shape, of the treatment area, known as "paradoxical hyperplasia", in the months following the treatment. If such paradoxical hyperplasia occurs, it will be distinguishable from temporary swelling and will probably not resolve on its own. The enlargement/lump can be removed by means of a surgical procedure such as liposuction Initial:
»Treatment area demarcationA small number of patients have experienced excessive fat removal in the treatment area, resulting in an unwanted indentation. The indentation may be improved through corrective procedures. Initial:
» In rare cases, patients have reported the CoolSculpting treatment area to have darker skin color, hardness, discrete nodules, frostbite (local injury due to cold), hernia or worsening of pre-existing hernia. Surgical intervention may be required correct hernia formation. Initial:
» Patient experiences may vary. Some patients may experience a delayed onset of the previously mentioned symptoms. Contact your physician immediately if any unusual side effects occur or if symptoms worsen over time. Initial:
»I understand that these and other unknown side effects may also occur. Initial:
Results Results
»You may start to see changes in as early as three weeks after your CoolSculpting procedure, and will experience the most dramatic results after one to three months. Your body will continue to naturally process the injured fat cells from your body for approximately four months after your procedure. Initial:
»Results vary from person to person. You may decide that additional treatments are necessary to achieve your desired outcome. Although highly unlikely, it is possible that you will not experience any noticeable result from the procedure. Initial:
Do you currently have or have had any of the following?
» Cryoglobulinemia (a condition in which an abnormal level of proteins thicken the blood in cold temperatures), or paroxysm cold hemoglobinuria or cold agglutinin disease (blood disorders in which cold temperatures lead to red blood cell death). Yes / No
» Known sensitivity to cold such as cold urticaria (hives triggered by cold), Raynaud's disease (disorder in which cold leads

Page 2 of 3

reduced blood flow in the fingers, which appear bumps that occur as a reaction to cold)	, ·	
» Poor blood flow in the area to be treated		Yes / No
» Neuropathic (nerve) disorders such as post-he	erpetic neuralgia or diabetic neuropa	athyYes / No
» Impaired skin sensation		Yes / No
» Open or infected wounds		Yes / No
» Bleeding disorders or use of blood thinners		Yes / No
» Recent surgery or scar tissue in the area to be	e treated	Yes / No
» A hernia or history of hernia in the area to be t	reated or adjacent to treatment site	Yes / No
» Skin conditions such as eczema, dermatitis, o	r rashes	Yes / No
» Pregnancy or lactation (making breast milk or	breast feeding)	Yes / No
» Any active implanted devices such as pacema	akers and defibrillators	Yes / No
» Any major health problems such as liver disea	ase	Yes / No
» Any known sensitivity to isopropyl alcohol (rub	bing alcohol) or propylene glycol	Yes / No
Pictures will be obtained for medical records. If pictures will be cropped or removed. (See Patient Photo		
As with most medical procedures, there are risk above information, and I give my informed corthis practice and his/her designated staff.		•
Print Name:S	ignature:	Date:
Witness:	Date:	
Physician(s):	Practice Name:	